arXiv:1507.02536v2 [math.CO] 22 Jul 2015

CORRIGENDUM
‘On the signless Laplacian spectra of k-trees’– CORRIGENDUM*

Minjie Zhang[a,b,†], Shuchao Li[a]

[a]Faculty of Mathematics and Statistics, Central China Normal University, Wuhan 430079, P.R. China
[b]Faculty of Mathematics and Physics, Hubei Institute of Technology, Huangshi 435003, P.R. China

A *k-tree* is either a complete graph on k vertices or a graph obtained from a smaller k-tree by adjoining a new vertex together with k edges connecting it to a k-clique. We tried to determine the n-vertex k-trees with the first three largest signless Laplacian indices in [Linear Algebra Appl. 467 (2015) 136-148]. Our proofs were mainly based on Lemma 2.4 in the above paper, whereas Lemma 2.4 follows from the next lemma:
(•) (Lemma 2.3 of [2]) *Let $0 < a_i \leqslant 1, 0 < b_i \leqslant 1$ for $1 \leqslant i \leqslant k$. Then $\sum_{i=1}^k b_i^2 - \sum_{i=1}^k a_i^2 \leqslant 2\left(\sum_{i=1}^k b_i - \sum_{i=1}^k a_i\right)$ and the equality holds if and only if $a_i = b_i = 1$ for $i = 1, 2, \ldots, k$.*

In fact, in the proof of Lemma 2.3, if $b_i - a_i < 0$, one cannot conclude the result. For example if $\sum_1^k b_i = \sum_1^k a_i$, we cannot conclude $\sum_1^k b_i^2 - \sum_1^k a_i^2 \leqslant 0$. Hence, Lemma 2.3 is not correct, which implies that our method in [2] can not be used to determine the n-vertex k-trees with the first three largest signless Laplacian indices.

The text below give a new method to determine the n-vertex k-trees with the first three largest signless Laplacian indices.

1. Preliminaries

We only consider simple connected graphs $G = (V_G, E_G)$, where V_G is the vertex set and E_G is the edge set. We call $n = |V_G|$ the *order* of G and $m = |E_G|$ the *size* of G. For a vertex subset S of V_G, denoted by $G[S]$ the subgraph induced by S. Let $N_G(v)$ denote the set of vertices adjacent to v. $N_G[v] = N_G(v) \cup \{v\}$. Let $A(G)$ be the adjacency matrix of graph G, $D(G)$ be the diagonal matrix with degrees of the vertices on the main diagonal. The signless matrix $Q(G) = D(G) + A(G)$ is real symmetric, its eigenvalues can be arranged as $q_1(G) \geqslant q_2(G) \geqslant \cdots \geqslant q_n(G) \geqslant 0$, where $q_1(G)$ is the *signless Laplacian index* of graph G.



$S_{k,n-k}$ G_1

Figure 1: Graphs $S_{k,n-k}$ and G_1.

For positive integers n, k with $n \geqslant k$, the *k-star* $S_{k,n-k}$ is depicted in Fig. 1. Let G_1 be the graph obtained from $S_{k,n-k}$ by deleting the edge u_1v_k and adding an edge u_1u_2 (see Fig. 1). For convenience, let $\mathscr{T}_n^k$ denote the

---

*Financially supported by the National Natural Science Foundation of China (Grant Nos. 11271149,11371162), the Program for New Century Excellent Talents in University (Grant No. NCET-13-0817) and the Special Fund for Basic Scientific Research of Central Colleges (Grant No. CCNU13F020).

†Corresponding author.

E-mail: zmj1982@21cn.com(M.J. Zhang), lscmath@mail.ccnu.edu.cn (S.C. Li)

set of all k-trees on n vertices. Let T_n^k be an n-vertex k-tree. Obviously, $T_n^k \cong K_k$ if $n = k$. So we consider T_n^k with $n \geqslant k+1$ in the whole context. A vertex $v \in V_{T_n^k}$ is called a *k-simplicial vertex* if v is a vertex of degree k whose neighbors form a k-clique of T_n^k. Let $S_1(T_n^k)$ be the set of all the k-simplicial vertices of T_n^k and we can easily have the following facts:

Fact 1. *$S_1(T_n^k) = \emptyset$ if and only if $T_n^k \cong K_k$; For any v in $V_{K_{k+1}}$, v is in $S_1(K_{k+1})$; If $n \geqslant k+2$, then $|S_1(T_n^k)| \geqslant 2$ and $S_1(T_n^k)$ is an independent set.*

Fact 2. *Let T_n^k be a k-tree on $n \geqslant k+2$ vertices. Then $d_{T_n^k}(v) = k$ for any $v \in S_1(T_n^k)$ and $T_n^k - v$ is also a k-tree.*

Fact 3. *$|S_1(T_n^k)| = n-k$ if and only if $T_n^k \cong S_{k,n-k}$.*

Given a k-clique $G[\{v_1, v_2, \dots, v_k\}]$, if there exists another vertex w in G such that $G[\{v_1, v_2, \dots, v_k, w\}]$ is a $(k+1)$-clique and $d_G(w) = k$, then we say that $G[\{v_1, v_2, \dots, v_k, w\}]$ has the *property* $P_G(v_1, v_2, \dots, v_k)$. Let $l_G(v_1, v_2, \dots, v_k)$ be the number of $(k+1)$-cliques which has the property $P_G(v_1, v_2, \dots, v_k)$ and set

$$l(G) := \max_{\{v_1,\dots,v_k\}\subseteq V_G} \{l_G(v_1, v_2, \dots, v_k) : G[\{v_1, v_2, \dots, v_k\}] \text{ is a } k\text{-clique of } G\}.$$

Obviously, $l(G) = n-k$ if and only if $G \cong S_{k,n-k}$, whereas $l(G) = n-k-2$ if and only if $G \cong G_1$; and there does not exist an n-vertex k-tree, say G, with $l(G) = n-k-1$.

Fact 4. *Let G be an n-vertex k-tree. Then $l(G) = n-k-3$ if and only if $G \cong G_2, G_3, G_4$ or G_5, where $G_2 = G_1 - u_3v_k + u_3u_2$, $G_3 = G_1 - u_3v_k + u_3u_1$, $G_4 = G_1 - u_3v_1 + u_3u_2$, $G_5 = G_1 - u_3v_{k-1} - u_3v_k + u_3u_1 + u_3u_2$. Graphs G_2, G_3, G_4, G_5 are depicted in Fig. 2.*

Fact 5. *For graphs G_2, G_3, G_4, G_5 (see Fig. 2), one has $|S_1(G_2)| = |S_1(G_4)| = n-k-1$, $|S_1(G_3)| = |S_1(G_5)| = n-k-2$.*



Figure 2: Graphs G_2, G_3, G_4 and G_5.

Further on we will need the following lemma.

Lemma 1.1 ([1])**.** *Let u and v be two distinct vertices of a connected graph G. Suppose that $w_1, w_2, \dots, w_s$ $(s \geqslant 1)$ are neighbors of v but not u and they are all different from u. Let $\mathbf{x} = (x_1, x_2, \dots, x_n)^T$ be the Perron vector of $Q(G)$, and let G^* be obtained from G by deleting the edges vw_i and adding the edges uw_i for $i = 1, 2, \dots, s$. If $x_v \leqslant x_u$, then $q_1(G) < q_1(G^*)$.*

2. Characterization of the k-trees with the first three largest signless Laplacian indices

In this section, we characterize the k-trees on n vertices having the maximal, second maximal and third maximal signless Laplacian indices.

Lemma 2.1. *Let G be in $\mathscr{T}_n^k \setminus \{S_{k,n-k}\}$ on $n \geqslant k+1$ vertices. Then there exists a k-tree G^* on n vertices such that $q_1(G) < q_1(G^*)$ with $|S_1(G^*)| = |S_1(G)| + 1$.*

Proof. Note that $G \ncong S_{k,n-k}$. By Fact 3, we get $|S_1(G)| < n-k$. Then, $G - S_1(G)$ is a k-tree whose order is no less than $k+1$ and $S_1(G - S_1(G)) \neq \emptyset$ by Facts 1 and 2. Let u be a vertex in $S_1(G - S_1(G))$ with $N_{G-S_1(G)}(u) = \{v_1, v_2 \ldots, v_k\}$. Assume that $N_G(u) \cap S_1(G) = \{w_1, w_2, \ldots, w_s\}$. Then, $d_G(w_j) = k$ and $N_G(w_j) \subseteq \{v_1, v_2, \ldots, v_k, u\}$ for any $1 \leqslant j \leqslant s$ by the definition of the k-tree. In order to complete the proof, it suffices to show the following two claims.

Claim 1. *If G contains two vertices w_i and w_j in $\{w_1, w_2, \ldots, w_s\}$ satisfying $N_G(w_i) \neq N_G(w_j)$, then there exists an n-vertex k-tree G' such that $q_1(G) < q_1(G')$ and $N_{G'}(w_i) = N_{G'}(w_j)$ with $|S_1(G')| = |S_1(G)|$ or $|S_1(G)| + 1$.*

Proof of Claim 1. Without loss of generality, we assume $N_G(w_i) = \{v_1, \ldots, v_k, u\} \setminus \{v_i\}$, $N_G(w_j) = \{v_1, \ldots, v_k, u\} \setminus \{v_j\}$ and

$$l_G(v_1, \ldots, v_{i-1}, v_{i+1}, \ldots, v_k, u) = a, \qquad l_G(v_1, \ldots, v_{j-1}, v_{j+1}, \ldots, v_k, u) = b,$$

where $v_i \neq v_j$. Hence, $1 \leqslant a \leqslant s$, $1 \leqslant b \leqslant s$. Let $\delta_1, \delta_2, \ldots, \delta_a$ (resp. $\Delta_1, \Delta_2, \ldots, \Delta_b$) be the $(k+1)$-cliques with the property $P_G(v_1, \ldots, v_{i-1}, v_{i+1}, \ldots, v_k, u)$ (resp. $P_G(v_1, \ldots, v_{j-1}, v_{j+1}, \ldots, v_k, u)$) and $w_{i_1}(= w_i), w_{i_2}, \ldots, w_{i_a}$ (resp. $w_{j_1}(= w_j), w_{j_2}, \ldots, w_{j_b}$) be the vertices in these $(k+1)$-cliques different from $v_1, \ldots, v_{i-1}, v_{i+1}, \ldots, v_k, u$ (resp. $v_1, \ldots, v_{j-1}, v_{j+1}, \ldots, v_k, u$). Suppose, without loss of generality, that $x_{v_i} \geqslant x_{v_j}$. Let

$$G' = G - \{v_j w_{i_1}, v_j w_{i_2}, \ldots, v_j w_{i_a}\} + \{v_i w_{i_1}, v_i w_{i_2}, \ldots, v_i w_{i_a}\}.$$

Then we have $G' \in \mathscr{T}_n^k$ with $N_{G'}(w_i) = N_{G'}(w_j)$ and $q_1(G) < q_1(G')$ by Lemma 1.1. Note that

$$\{v_1, \ldots, v_{j-1}, v_{j+1}, \ldots, v_k, u, w_{i_1}, w_{i_2}, \ldots, w_{i_a}\} \subseteq N_G(v_j).$$

Hence, we get $S_1(G') = S_1(G)$ if $d_G(v_j) > k+a$; whereas $S_1(G') = S_1(G) \cup \{v_j\}$ if $d_G(v_j) = k+a$. Thus, $|S_1(G')| = |S_1(G)|$ or $|S_1(G)| + 1$, as desired. □

Claim 2. *If $N_G(w_1) = N_G(w_2) = \cdots = N_G(w_s)$, then there exists an n-vertex k-tree G^* such that $q_1(G) < q_1(G^*)$ and $|S_1(G^*)| = |S_1(G)| + 1$.*

Proof of Claim 2. Without loss of generality, we assume $N_G(w_j) = \{v_1, \ldots, v_{k-1}, u\}$ for $1 \leqslant j \leqslant s$. Let $V_0 = N_G(v_k) \setminus \{v_1, \ldots, v_{k-1}, u\}$. Then, $V_0 \neq \emptyset$ since $v_k \notin S_1(G)$. If $x_{v_k} \geqslant x_u$, then let

$$G^* = G - \{uw_1, uw_2, \ldots, uw_s\} + \{v_k w_1, v_k w_2, \ldots, v_k w_s\}.$$

It is routine to check that G^* is an n-vertex k-tree and $|S_1(G^*)| = |S_1(G) \cup \{u\}| = |S_1(G)| + 1$. By Lemma 1.1, $q_1(G) < q_1(G^*)$.

If $x_{v_k} < x_u$, then let $G^* = G - \sum_{v \in V_0} v_k v + \sum_{v \in V_0} uv$. It is routine to check G^* is an n-vertex k-tree and $|S_1(G^*)| = |S_1(G) \cup \{v_k\}| = |S_1(G)| + 1$. By Lemma 1.1, $q_1(G) < q_1(G^*)$.

This completes the proof of Claim 2. □

Now we come back to the proof of Lemma 2.1. If $N_G(w_1) = N_G(w_2) = \cdots = N_G(w_s)$, by Claim 2 our result holds. Otherwise, repeatedly using Claim 1 (at most $(s-1)$ times), one can finally get an n-vertex k-tree G'' with $q_1(G) < q_1(G'')$ satisfying one of the following:

(i) $|S_1(G'')| = |S_1(G)| + 1$;
(ii) $|S_1(G'')| = |S_1(G)|$ and $N_{G''}(w_1) = N_{G''}(w_2) = \cdots = N_{G''}(w_s)$.

If (i) holds, then let $G^* \cong G''$, our lemma holds obviously. If (ii) holds, combining with Claim 2, we can get a k-tree G^* on n vertices such that $q_1(G'') < q_1(G^*)$ and $|S_1(G^*)| = |S_1(G'')| + 1 = |S_1(G)| + 1$, as desired. $\square$

The following theorem follows immediately from Lemma 2.1.

Theorem 2.2. *Let G be an n-vertex k-tree with $n \geqslant k+1$. Then $q_1(G) \leqslant q_1(S_{k,n-k})$ and the equality holds if and only if $G \cong S_{k,n-k}$, where $S_{k,n-k}$ is depicted in Fig. 1.*

Lemma 2.3. *Let G be an n-vertex k-tree with $n \geqslant k+1$ and $|S_1(G)| = n-k-1$. If $G \ncong G_1$, then there exists an n-vertex k-tree G^* with $|S_1(G^*)| = n-k-1$ such that $q_1(G) < q_1(G^*)$ and $l(G^*) = l(G) + 1$.*

Proof. Note that $|S_1(G)| = n-k-1$. Hence, the order of $G - S_1(G)$ is $k+1$. By Fact 2, $G - S_1(G)$ is a k-tree, that is, $G - S_1(G) \cong K_{k+1}$. Denote the vertex set of $G - S_1(G)$ by $\{v_1, v_2, \ldots, v_{k+1}\}$.

As $G \ncong G_1$, we have $l(G) < n-k-2$. Assume, without loss of generality, that $l(G) = l_G(v_1, v_2, \ldots, v_k) = s$. Hence, $s < n-k-2$. Let $\delta_1, \delta_2, \ldots, \delta_s$ be the $(k+1)$-cliques satisfying the property $P_G(v_1, v_2, \ldots, v_k)$ and $w_1, w_2, \ldots, w_s$ be the vertices in these $(k+1)$-cliques different from $v_1, v_2, \ldots, v_k$. Obviously, $d_G(w_1) = d_G(w_2) = \cdots = d_G(w_s) = k$ and $w_iw_j \notin E_G$ for $1 \leqslant i < j \leqslant s$.

Note that $|S_1(G)| = n-k-1$ and $s < n-k-2$. Hence, there exists a vertex $r_1 \notin \{w_1, w_2, \ldots, w_s\}$ such that $d_G(r_1) = k$ and $G[N_G[r_1]] \cong K_{k+1}$. Without loss of generality, we assume $N_G(r_1) = \{v_2, v_3, \ldots, v_{k+1}\}$ and $l_G(v_2, v_3, \ldots, v_{k+1}) = t$. Obviously, $1 \leqslant t \leqslant s$. Let $\Delta_1, \Delta_2, \ldots, \Delta_t$ be the $(k+1)$-cliques satisfying the property $P_G(v_2, v_3, \ldots, v_{k+1})$ and $r_1, r_2, \ldots, r_t$ be the vertices in these $(k+1)$-cliques different from $v_2, v_3, \ldots, v_{k+1}$. Suppose that $\mathbf{x} = (x_1, x_2, \ldots, x_n)^T$ is the Perron vector of $Q(G)$.

If $x_{v_1} \geqslant x_{v_{k+1}}$, then let $G^* = G - v_{k+1}r_1 + v_1r_1$. It is routine to check that $G^* \in \mathscr{T}_n^k$. By Lemma 1.1, $q_1(G) < q_1(G^*)$. As $s+1 < n-k-1 = |S_1(G)|$, there exists a vertex $y \in S_1(G) \setminus \{w_1, w_2, \ldots, w_s, r_1\}$. Note that $G[N_G(y)]$ is a k-clique not satisfying the property $P_G(v_1, v_2, \ldots, v_k)$. Hence, $yv_{k+1} \in E_G$. Thus, $v_{k+1} \notin S_1(G^*)$. We have $l(G^*) = |\{w_1, w_2, \ldots, w_s, r_1\}| = l(G) + 1$ and $S_1(G^*) = S_1(G)$, as desired.

If $x_{v_1} < x_{v_{k+1}}$, then let

$$G^* = G - \{v_1w_1, v_1w_2, \ldots, v_1w_{s-t+1}\} + \{v_{k+1}w_1, v_{k+1}w_2, \ldots, v_{k+1}w_{s-t+1}\}.$$

Then G^* is in $\mathscr{T}_n^k$ and by Lemma 1.1, $q_1(G) < q_1(G^*)$. Note that if $t \geqslant 2$, then one may easily get $v_1 \notin S_1(G^*)$. Now we consider that $t = 1$. In this case, as $s+1 < n-k-1 = |S_1(G)|$, there exists a vertex $z \in S_1(G) \setminus \{w_1, w_2, \ldots, w_s, r_1\}$. Note that $G[N_G(z)]$ is a k-clique not satisfying the property $P_G(v_2, v_3, \ldots, v_{k+1})$. Hence, $zv_1 \in E_G$ and $v_1 \notin S_1(G^*)$. Thus, $l(G^*) = |\{r_1, r_2, \ldots, r_t, w_1, w_2, \ldots, w_{s-t+1}\}| = s+1 = l(G)+1$ and $S_1(G^*) = S_1(G)$, as desired. $\square$

The following theorem follows immediately from Lemmas 2.1 and 2.3.

Theorem 2.4. *Let G be in $\mathscr{T}_n^k \setminus \{S_{k,n-k}\}$ with $n \geqslant k+1$. Then $q_1(G) \leqslant q_1(G_1)$ with equality if and only if $G \cong G_1$, where G_1 is depicted in Fig. 1.*

Lemma 2.5. *Let G be an n-vertex k-tree with $n \geqslant k+1$ and $l(G) = n-k-3$. Then $q_1(G) \leqslant q_1(G_2)$ with equality if and only if $G \cong G_2$, where G_2 is depicted in Fig. 2.*

Proof. As G is an n-vertex k-tree with $l(G) = n-k-3$, by Fact 4, it suffices to show the next three claims.

Claim 1. $q_1(G_3) < q_1(G_2)$.

Proof of Claim 1. Let $\mathbf{x} = (x_1, x_2, \ldots, x_n)^T$ be the Perron vector of $Q(G_3)$. Put $G' := G_3 - u_1u_3 + u_2u_3$ if $x_{u_1} \leqslant x_{u_2}$ and $G'' := G_3 - u_2v_k + u_1v_k$ otherwise. It is routine to check that $G' \cong G'' \cong G_2$. By Lemma 1.1, we have $q_1(G_3) < q_1(G_2)$, as desired. $\square$

By the same discussion as in the proof of Claim 1 as above, we can show the next claim, which is omitted.

Claim 2. $q_1(G_4) < q_1(G_2)$.

Claim 3. $q_1(G_5) < q_1(G_2)$.

Proof of Claim 3. Let $\mathbf{x} = (x_1, x_2, \ldots, x_n)^T$ be the Perron vector of $Q(G_5)$. If $x_{v_{k-1}} \geqslant x_{u_2}$, then let $G' := G_5 - u_2u_3 + v_{k-1}u_3$. Clearly, $G' \cong G_3$. By Lemma 1.1 and Claim 1, one has $q_1(G_5) < q_1(G_3) < q_1(G_2)$, as desired.

Now we consider $x_{v_{k-1}} < x_{u_2}$. For convenience, let $V_0 = N_{G_5}(v_{k-1}) \setminus \{v_1, v_2, \ldots, v_{k-2}, v_k, u_1, u_2\}$. In this case, put $G'' := G_5 - \sum_{v \in V_0} vv_{k-1} + \sum_{v \in V_0} vu_2$. It is routine to check that $G'' \cong G_3$. By Lemma 1.1 and Claim 1, one has $q_1(G_5) < q_1(G_3) < q_1(G_2)$, as desired. $\square$

By Claims 1-3, Lemma 2.5 holds. $\square$

Lemma 2.6. *Let G be an n-vertex k tree with $n \geqslant k+1$ and $|S_1(G)| = n-k-2$. If $l(G) < n-k-3$, then there exists an n-vertex k tree G^* with $l(G^*) = l(G)+1$ such that $q_1(G) < q_1(G^*)$ and one of the following holds:*

$$|S_1(G^*)| = n-k-2, \tag{2.1}$$

$$|S_1(G^*)| = n-k-1 \text{ and } G^* \cong G_2. \tag{2.2}$$

Proof. Note that $|S_1(G)| = n-k-2$; hence, together with Fact 2, $G - S_1(G)$ is a $(k+2)$-vertex k-tree, i.e., $G - S_1(G) \cong S_{k,2}$. Let $V_{G-S_1(G)} = \{v_1, v_2, \ldots, v_{k+2}\}$ with $d_{G-S_1(G)}(v_{k+1}) = d_{G-S_1(G)}(v_{k+2}) = k$. Obviously, $v_{k+1}v_{k+2} \notin E_G$. Clearly, $l(G) \geqslant l_G(v_1, v_2, \ldots, v_k)$. Put $l_G(v_1, v_2, \ldots, v_k) = s$.

Case 1. $l(G) = l_G(v_1, v_2, \ldots, v_k)$. In this case, one has $s < n-k-3$. Let $\delta_1, \delta_2, \ldots, \delta_s$ be the $(k+1)$-cliques satisfying the property $P_G(v_1, v_2, \ldots, v_k)$ and $w_1, w_2, \ldots, w_s$ be the vertices in these $(k+1)$-cliques different from $v_1, v_2, \ldots, v_k$. Then by Fact 1, $d_G(w_1) = d_G(w_2) = \cdots = d_G(w_s) = k$ and $w_iw_j \notin E_G$ for $1 \leqslant i < j \leqslant s$.

Obviously, there exists a vertex r_1 in $S_1(G)$ such that $v_{k+1}r_1 \in E_G$ by the definitions of the k-tree and $S_1(G)$. Assume, without loss of generality, that $N_G(r_1) = \{v_2, v_3, \ldots, v_k, v_{k+1}\}$ and $l_G(v_2, v_3, \ldots, v_k, v_{k+1}) = t$. Then, $1 \leqslant t \leqslant s$. Let $\Delta_1, \Delta_2, \ldots, \Delta_t$ be the $(k+1)$-cliques satisfying the property $P_G(v_2, v_3, \ldots, v_k, v_{k+1})$ and $r_1, r_2, \ldots, r_t$ be the vertices in these $(k+1)$-cliques different from $v_2, v_3, \ldots, v_k, v_{k+1}$. Suppose that $\mathbf{x} = (x_1, x_2, \ldots, x_n)^T$ is the Perron vector of $Q(G)$.

Since $v_{k+1}, v_{k+2} \notin S_1(G)$, we get $d_G(v_{k+1}), d_G(v_{k+2}) \geqslant k+1$. We proceed by considering the following two possible subcases:

Subcase 1.1. There exists a vertex, say v_{k+1}, in $\{v_{k+1}, v_{k+2}\}$ such that $d_G(v_{k+1}) > k+1$. If $x_{v_1} \geqslant x_{v_{k+1}}$, then let $G^* = G - v_{k+1}r_1 + v_1r_1$. By Lemma 1.1, we have $q_1(G) < q_1(G^*)$. Combining with $d_G(v_{k+1}) > k+1$, one can easily check that $l(G^*) = l(G)+1$ and $S_1(G^*) = S_1(G)$. If $x_{v_1} < x_{v_{k+1}}$, then let

$$G^* = G - \{v_1w_1, v_1w_2, \ldots, v_1w_{s-t+1}\} + \{v_{k+1}w_1, v_{k+1}w_2, \ldots, v_{k+1}w_{s-t+1}\}.$$

Then G^* is in $\mathscr{T}_n^k$ and by Lemma 1.1, $q_1(G) < q_1(G^*)$. Note that $\{v_2, v_3, \ldots, v_k, v_{k+1}, v_{k+2}\} \subseteq N_{G^*}(v_1)$, we also get $l(G^*) = l(G)+1$ and $S_1(G^*) = S_1(G)$, our Lemma holds in this case.

Subcase 1.2. $d_G(v_{k+1}) = d_G(v_{k+2}) = k+1$. In this case, we have $s = l(G) = n-k-4$. Without loss of generality, we assume $x_{v_{k+1}} \leqslant x_{v_{k+2}}$ and let $G' = G - v_{k+1}r_1 + v_{k+2}r_1$. By Lemma 1.1, we have $q_1(G) < q_1(G')$. Combining with $d_G(v_{k+1}) = k+1$, we get

$$l(G') = |\{w_1, w_2, \ldots, w_s, v_{k+1}\}| = l(G)+1 = n-k-3.$$

However, noting that $S_1(G') = S_1(G) \cup \{v_{k+1}\}$, hence, $|S_1(G')| = |S_1(G)| + 1 = n - k - 1$. By Facts 4 and 5, $G' \cong G_2$ or G_4. Note that $q_1(G_4) < q_1(G_2)$; hence by Lemma 2.5, our lemma holds in this case.

Case 2. $l(G) > l_G(v_1, v_2, \dots, v_k)$. Without loss of generality, we assume that $l(G) = l_G(v_2, v_3, \dots, v_k, v_{k+1}) = s$. Hence, $s < n - k - 3$. Let $\delta_1, \delta_2, \dots, \delta_s$ be the $(k+1)$-cliques satisfying the property $P_G(v_2, v_3, \dots, v_k, v_{k+1})$ and $w_1, w_2, \dots, w_s$ be the vertices in these $(k+1)$-cliques different from $v_2, v_3, \dots, v_k, v_{k+1}$.

If $l_G(v_1, v_2, \dots, v_k) \neq 0$, then by a similar discussion as in the proof of Case 1, one can show this lemma is true. Hence, we consider that $l_G(v_1, v_2, \dots, v_k) = 0$. By the definitions of the k-tree and $S_1(G)$, it is routine to check that there exists a vertex r_1 in $S_1(G)$ such that $v_{k+2}r_1 \in E_G$. Thus we have $v_{k+1}r_1 \notin E_G$ and there exists only one vertex, say v_i, in $\{v_1, v_2, \dots, v_k\}$ such that $v_ir_1 \notin E_G$. If $i = 1$, then $N_G(r_1) = \{v_2, v_3, \dots, v_k, v_{k+2}\}$. By a similar discussion as in the proof of Case 1, one can show this lemma holds. In what follows we consider that $i \neq 1$. Then, $N_G(r_1) = \{v_1, v_2, \dots, v_{i-1}, v_{i+1}, \dots, v_k, v_{k+2}\}$.

Assume that $l_G(v_1, v_2, \dots, v_{i-1}, v_{i+1}, \dots, v_k, v_{k+2}) = t$, then $1 \leqslant t \leqslant s$. Let $\Delta_1, \Delta_2, \dots, \Delta_t$ be the $(k+1)$-cliques satisfying the property $P_G(v_1, v_2, \dots, v_{i-1}, v_{i+1}, \dots, v_k, v_{k+2})$ and $r_1, r_2, \dots, r_t$ be the vertices in these $(k+1)$-cliques different from $v_1, v_2, \dots, v_{i-1}, v_{i+1}, \dots, v_k, v_{k+2}$. We proceed through the following two possible subcases.

Subcase 2.1. $x_{v_1} \leqslant x_{v_i}$. Then let $G' = G - v_1r_1 + v_ir_1$. By Lemma 1.1, we have $q_1(G) < q_1(G')$. Note that $\{v_2, v_3, \dots, v_k, v_{k+1}, v_{k+2}\} \subseteq N_{G'}(v_1)$; hence we get $l(G') = l(G), S_1(G') = S_1(G)$ and $N_{G'}(r_1) = \{v_2, v_3, \dots, v_k, v_{k+2}\}$. By a similar discussion as in the proof of Case 1, one can find an n-vertex k-tree G^* with $l(G^*) = l(G') + 1$ such that $q_1(G') < q_1(G^*)$ and either (2.1) or (2.2) holds. Hence, we have $q_1(G) < q_1(G') < q_1(G^*)$ and $l(G^*) = l(G') + 1 = l(G) + 1$, as desired.

Subcase 2.2. $x_{v_1} > x_{v_i}$. Obviously, $G\ [\{v_1, v_2, \dots, v_{i-1}, v_{i+1}, \dots, v_k, v_{k+1}\}\ ]$ is a k-clique. Assume that $l_G(v_1, v_2,$
$\dots, v_{i-1}, v_{i+1}, \dots, v_k, v_{k+1}) = a$ where $0 \leqslant a \leqslant s$. Let $\Delta'_1, \Delta'_2, \dots, \Delta'_a$ be the $(k+1)$-cliques satisfying the property $P_G(v_1, v_2, \dots, v_{i-1}, v_{i+1}, \dots, v_k, v_{k+1})$ and $y_1, y_2, \dots, y_a$ be the vertices in these $(k+1)$-cliques different from $v_1, v_2, \dots, v_{i-1}, v_{i+1}, \dots, v_k, v_{k+1}$. Put

$$G'' := G - \{v_iw_1, v_iw_2, \dots, v_iw_{s-a}\} + \{v_1w_1, v_1w_2, \dots, v_1w_{s-a}\}.$$

Then G'' is in $\mathscr{T}_n^k$ and by Lemma 1.1 we have $q_1(G) < q_1(G'')$. Note that $\{v_1, \dots, v_{i-1}, v_{i+1}, \dots, v_k, v_{k+1}, v_{k+2}\} \subseteq N_{G''}(v_i)$; hence we get $l(G'') = l(G), S_1(G'') = S_1(G)$. On the other hand, we have

$$N_{G''}(y_t) = N_{G''}(w_j) = \{v_1, v_2, \dots, v_{i-1}, v_{i+1}, \dots, v_k, v_{k+1}\}$$

for all $0 \leqslant t \leqslant a$ and $1 \leqslant j \leqslant s - a$. By a similar discussion as in the proof of Case 1, there exists an n-vertex k-tree G^* with $l(G^*) = l(G'') + 1$ such that $q_1(G'') < q_1(G^*)$ and either (2.1) or (2.2) holds. Hence, we have $q_1(G) < q_1(G'') < q_1(G^*)$ and $l(G^*) = l(G'') + 1 = l(G) + 1$, as desired.

This completes the proof. □

Theorem 2.7. *Let G be in $\mathscr{T}_n^k \setminus \{S_{k,n-k}, G_1\}$ on $n \geqslant k + 1$ vertices. Then, $q_1(G) \leqslant q_1(G_2)$ and the equality holds if and only if $G \cong G_2$, where G_2 is depicted in Fig. 2.*

Proof. Since $G \ncong S_{k,n-k},\ G_1$, we have $|S_1(G)| \leqslant n-k-1$ and $l(G) \leqslant n-k-3$. Note that $|S_1(G_1)| = |S_1(G_2)| = n - k - 1$; hence we proceed by considering the following possible cases.

Case 1. $|S_1(G)| = n - k - 1$. In this case, let G be in $\mathscr{T}_n^k \setminus \{G_1\}$ with $|S_1(G)| = n - k - 1$ such that $q_1(G)$ is as large as possible. By Lemma 2.3, we have $l(G) = n - k - 3$. By Facts 4 and 5 we have $G \cong G_2$ or G_4. In view of Lemma 2.5, $q_1(G_4) < q_1(G_2)$, one can easily get $q_1(G) \leqslant q_1(G_2)$ when $G \in \mathscr{T}_n^k \setminus \{G_1\}$ and the equality holds if and only if $G \cong G_2$.

Case 2. $|S_1(G)| = n-k-2$. In this case, by Fact 2 we obtain that $G - S_1(G)$ is a k-tree on $k+2$ vertices, that is, $G - S_1(G) \cong S_{k,2}$. Denote the vertex set of $G - S_1(G)$ by $\{v_1, v_2, \ldots, v_{k+2}\}$ with $d_{G-S_1(G)}(v_{k+1}) = d_{G-S_1(G)}(v_{k+2}) = k$. Obviously, $v_{k+1}v_{k+2} \notin E_G$ and there exist vertices w_1 and r_1 in $S_1(G)$ such that $v_{k+1}w_1$, $v_{k+2}r_1 \in E_G$ by the definitions of the k-tree and $S_1(G)$. Thus, $N_G(r_1) \neq N_G(w_1)$. Hence, $l(G) < |S_1(G)| = n-k-2$, i.e., $l(G) \leqslant n-k-3$.

Let G be in $\mathscr{T}_n^k$ with $|S_1(G)| = n-k-2$ such that $q_1(G)$ is as large as possible. By Lemma 2.6, one has either $l(G) = n-k-3$ or $q_1(G) < q_1(G_2)$. However, if $l(G) = n-k-3$, then combining with $|S_1(G)| = n-k-2$ and by Facts 4 and 5, we have $G \cong G_3$ or G_5. By Lemma 2.5, $q_1(G_5) < q_1(G_3) < q_1(G_2)$. Hence, we obtain $q_1(G) < q_1(G_2)$ if $|S_1(G)| = n-k-2$.

Case 3. $|S_1(G)| \leqslant n-k-3$. In this case, repeatedly using Lemma 3.1 yields a k-tree G' on n vertices such that $q_1(G) < q_1(G')$ with $|S_1(G')| = n-k-2$. However, in view of Case 2, we get $q_1(G') < q_1(G_2)$. So, $q_1(G) < q_1(G') < q_1(G_2)$ if G is an n-vertex k-tree with $|S_1(G)| \leqslant n-k-3$.

By Cases 1-3, Theorem 2.7 holds. □

Acknowledgements

We thank Dr. Asghar Bahmani for drawing our attention to the mistake.

References

[1] Y. Hong, X.D. Zhang, Sharp upper and lower bounds for largest eigenvalue of the Laplacian matrices of trees, Discrete Math. 296 (2005) 187-197.

[2] M.J. Zhang, S.C. Li, On the signless Laplacian spectra of k-trees, Linear Algebra Appl. 467 (2015) 136-148.